

EASY LIQUIDS ZN-SUL

PRODUCT GUIDE



OVERVIEW

Zinc is an important micro-nutrient for all crops and placement is very important as it is immobile in the soil. When building soil zinc levels, it is important to treat a significant proportion of the root zone. An application of EASY Liquids ZN-SUL with a boom spray and incorporated by tillage mixes zinc through the cultivated layer helping zinc availability during crop development. EASY Liquids are manufactured at our regionally located plants with the highest quality standards.

ANALYSIS

N		P	K	S	Ca	Mg	B	Cu	Fe	Mn	Mo	Zn	SG	
Total N	Nitrate	Ammonium	Urea	Phosphorus	Potassium	Sulphur	Calcium	Magnesium	Boron	Copper	Iron	Manganese	Molybdenum	Zinc
						8.5						16.0	1.38	

All nutrient content is shown in w/v.

USAGE RECOMMENDATIONS

PREPLANT

Can be soil applied prior to planting through ground and aerial application equipment to build soil zinc levels. It is best incorporated into the soil by a tillage operation as zinc is immobile.

- All broadacre crops: Spray up to 60 L/ha pre-planting and incorporate into soil or band with preplant nitrogen. Common rates 10-20 L/ha on lighter textured soils and 20-30 L/ha for heavy clay soils. Water rate: 40-50 L/ha.

PLANTING / SEEDING

Apply up to 10 L/ha (1 m row spacing) or 40 L/ha (25cm row spacing) in furrow with the seed during planting.

FERTIGATION

Apply through irrigation systems at up to 30 L/ha. (Total - split over multiple applications).

For detailed fertiliser recommendations for your crops please contact your local EASY Liquids sales agronomist.

EASY LIQUIDS ZN-SUL

PRODUCT GUIDE



OVERVIEW

Zinc is an important micro-nutrient for all crops and placement is very important as it is immobile in the soil. When building soil zinc levels, it is important to treat a significant proportion of the root zone. An application of EASY Liquids ZN-SUL with a boom spray and incorporated by tillage mixes zinc through the cultivated layer helping zinc availability during crop development. EASY Liquids are manufactured at our regionally located plants with the highest quality standards.

ANALYSIS

N		P	K	S	Ca	Mg	B	Cu	Fe	Mn	Mo	Zn	SG		
Total N	Nitrate	Ammonium	Urea	Phosphorus	Potassium	Sulphur	Calcium	Magnesium	Boron	Copper	Iron	Manganese	Molybdenum	Zinc	
						8.5						16.0		1.38	

All nutrient content is shown in w/v.

FOLIAR APPLICATION

COTTON

Foliar apply 0.5-1.5 L/ha from 4 leaf to the start of flowering. Minimum water rate 50 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 2 L of N-FOL per 100 L spray solution will assist uptake.

GRAIN LEGUMES

Summer: Foliar apply 0.5-1.5 L/ha from 4 leaf to the start of flowering. Minimum water rate 50 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 2 L of N-FOL per 100L spray solution will assist uptake.

Winter: Foliar apply 0.5-1.5 L/ha from 4 leaf to the start of flowering. Minimum water rate 50 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 2 L of N-FOL per 100L spray solution will assist uptake.

VEGETABLE

Foliar apply 0.35 L/ha from 2-3 weeks after planting/transplanting and repeat as required. Minimum water rate 250 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 1 L of N-FOL per 100 L spray solution will assist uptake.

SORGHUM

Foliar apply 0.5-1.5 L/ha from 3 to 6 leaf growth stage. Minimum water rate 50 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 2 L of N-FOL per 100 L spray solution will assist uptake.

TREE CROP AND VINES

(Fruit, Citrus & Nuts)

Foliar apply 0.14 L/ha from leaf out to early fruit development. Minimum water rate 1000 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 0.2 L of N-FOL per 100L spray solution will assist uptake.

MAIZE

Foliar apply 0.5-1.5L/ha from 2 to 6 leaf growth stage. Minimum water rate 50 L/ha by ground. The addition of 100 mL/100L of WETTER 1000 Surfactant and 2L of N-FOL per 100 L spray solution will assist uptake.

WINTER CEREAL

Foliar apply 0.5-1.5 L/ha from 5 leaf to end of tillering (GS Z15 to GS Z30). Minimum water rate 50 L/ha by ground. The addition of 100 mL/100 L of WETTER 1000 Surfactant and 2 L of N-FOL per 100 L spray solution will assist uptake.

QUALITY LIQUID NUTRIENTS, MADE EASY



EASY Liquids

To find out how EASY Liquids
could be easier for you and
your farm, talk to us today:

Moree 02 6752 7509
Boundary Bend 02 6752 7509
Whitton 02 6955 2565
au.sales@incitecpivot.com.au



Disclaimer: whilst these recommendations are typically right for most cropping situations, this is intended as a helpful guide. You should consult with your agronomist or crop nutrition expert to ensure you have the right product and application rate for your crop. Farming situations and nutrition requirements of crops vary.